

Research Article

Cite this article: Ayala Guanga L, Astiz S, Dutan Sanango J, Samaniego JX, López Cabrera RX, Perea F, Hernandez-Fonseca HJ and Pesántez Pacheco JL (2022). Relationship between the concentration of anti-Müllerian hormone and antral follicle count in pasture-managed Holstein cows in the high tropics. *Journal of Dairy Research* **89**, 60–64. <https://doi.org/10.1017/S0022029922000140>

Received: 18 May 2020
Revised: 22 November 2021
Accepted: 29 November 2021
First published online: 22 February 2022

Keywords:

Anti-Müllerian hormone; antral follicle; biomarker; bovine; dairy cattle

Author for correspondence:

Susana Astiz, Email: astiz.susana@inia.es

Relationship between the concentration of anti-Müllerian hormone and antral follicle count in pasture-managed Holstein cows in the high tropics

Luis Ayala Guanga¹, Susana Astiz², Jorge Dutan Sanango¹,
Jorge Xavier Samaniego¹, Rodrigo Xavier López Cabrera⁴, Fernando Perea⁵,
Hugo J. Hernandez-Fonseca³ and José Luis Pesántez Pacheco¹

¹Faculty of Agricultural Sciences, School of Veterinary Medicine and Zootechnics, University of Cuenca, Avda. 12 de octubre y Diego de Tapia, 010220 Cuenca, Ecuador; ²Animal Reproduction Department (INIA-CSIC), INIA, Avda. Puerta de Hierro s/n, 28040 Madrid, Spain; ³Faculty of Veterinary Sciences, Universidad del Zulia, Avda. 16 Guajira, 4005 Maracaibo, Venezuela; ⁴Program in Agricultural Biotechnology, Universidad Técnica Particular de Loja, San Cayetano Alto s/n, 110150 Loja, Ecuador and ⁵Department of Agricultural Sciences, Universidad de Los Andes, 3101 Trujillo, Venezuela

Abstract

We evaluated the relationship between plasma levels of anti-Müllerian hormone (AMH) and the number of antral follicles at the restart of the follicular wave in crossbred Holstein cows reared under extensive grazing systems over 2500 m above sea level. The study included 140 cows from 15 farms that were in average at the 75.3 ± 2.10 d post partum. Animals were synchronized according to the following regime: day 0 = intravaginal progesterone releasing device (IPD) + estradiol benzoate (EB); day 7: withdrawal of IPD + prostaglandin; and day 8: EB, for restart of the follicular wave on day 11. On this day 11, antral follicle counts (AFCs) were made by transrectal ultrasound, and a plasma sample was taken for the determination of AMH. The mean AMH plasma level was 0.06 ± 0.03 ng/ml and the mean AFC was 17.26 ± 0.38 follicles. A strong positive linear correlation was found between these two variables ($r = 0.783$, $r = 0.613$, $P < 0.0001$). Cows were categorized according to AMH concentration as high (>0.09 ng/ml), intermediate (0.09 – 0.05 ng/ml) or low (<0.05 ng/ml). Cows with high AMH presented a higher AFC (25.0 ± 2.21 follicles) than those with low AMH (14.08 ± 2.68 follicles; $P < 0.001$). Our results suggest that the cut-off value of AMH = 0.09 ng/ml may be useful for selecting donors in multiple ovulation embryo transfer programs involving cows with these characteristics. Our data further suggest that AMH plasma concentration correlates with AFC and can be used as an endocrine biomarker of the number of antral follicles present at a given moment of the estrous cycle in crossbred Holstein cows raised at altitudes above 2500 m.

Reproductive biotechnologies can contribute to rapid genetic improvement of cattle. However, their efficacy depends on physiological characteristics of the female such as the antral follicle count (AFC), which can vary substantially among donors. AFC may be a good marker for selecting donors with the greatest potential to generate competent oocytes, which would increase the proportion of animals that respond well to reproductive biotechnologies (Alward and Bohlen, 2020). At present, donors with high AFC are selected based on transrectal ultrasound in the framework of multiple ovulation embryo transfer (MOET) programs for assisted reproduction in cattle. However, transrectal ultrasound requires trained personnel, expensive equipment and adequate facilities for the careful evaluation of the obtained images (Perry and Cushman, 2016). An effective alternative to the ultrasound technique is the quantification of anti-Müllerian hormone (AMH), specifically produced by granulosa cells of growing small follicles (Monniaux *et al.*, 2010). Levels of this hormone are directly related to AFC in *Bos taurus* cows (Souza *et al.*, 2015). However, the level of AMH taken at an unknown moment of the estrous cycle varies with animals, with high individual repeatability (Gobikrushanth *et al.*, 2017), preventing the standardized use of AMH concentration as an endocrine marker without previous evidence of the intensity of such relationship at a given moment of the estrous cycle. Moreover, whether this relationship also holds in grazing feeding systems in the high tropics is unclear.

The recommended cut-off for AMH levels to identify animals with high AFC differs across subspecies; for example, they are 0.70 ng/ml for *Bos indicus* (Brahman breed) and 0.25 ng/ml for *Bos taurus* (Angus breed; Jaques *et al.*, 2020). The levels of AMH considered low or high can differ between breeds of the same *B. taurus* subspecies, and even in the same breed. In the

© The Author(s), 2022. Published by Cambridge University Press on behalf of Hannah Dairy Research Foundation. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



CAMBRIDGE
UNIVERSITY PRESS

Holstein breed raised under intensive production systems (in this case, Florida, USA), concentrations over 0.631 ng/ml are considered high and those under 0.085 ng/ml are considered low (Ribeiro *et al.*, 2014). Furthermore, different studies of the same breed have suggested different cut-off values. This variability in recommended cut-off points may be because the relationship of AMH levels with AFC in crossbred Holstein cattle depends on management and feeding conditions. In particular, we are interested in how levels of AMH and their relationship with AFC differ between animals under extensive grazing production in high tropical climatic conditions (over 2500 m above sea level) and animals under the same production system in other environments. Several considerations suggest that altitude may influence cattle fertility, although this has not been demonstrated directly. High altitude can alter physiological functions of animals (Parraguez and Gonzalez, 2020). The hypobaric hypoxia resulting from living at high altitude causes oxidative stress and affects the hypothalamus, pituitary and ovarian axis, compromising folliculogenesis of the *corpus luteum* and expression of the genes encoding insulin-like growth factors (IGFs) I and II. This effect of hypobaric hypoxia, in turn, reduces the fertility of animals living at high altitude compared to those living at sea level (Parraguez and Gonzalez, 2020). Criollo or also named Creole cattle in the highlands of Ecuador show differences in ovarian physiology from those reared at sea level, and these differences could be linked to altitude (Ayala *et al.*, 2019). The agroecological conditions typical of the area and the nutritional management in high-altitude regions could alter the energy balance as well as the metabolic and endocrine activities of these Creole cattle, affecting reproductive performance.

The objective of the present study was to determine the relationship between plasma AMH concentration and the AFC, assessed during the restart of the follicular wave, in crossbred Holstein cows raised in the high tropics under an extensive grazing system. Based on our analysis, we aimed to propose cut-off values for AMH levels to select animals with high AFC for MOET protocols, currently being implemented in the frame of official breed recovery programs.

Material and methods

Animals and farms

From a total of 609 farms located in the Ecuadorian high tropics, 15 farms were chosen based on altitude (over 2500 m above sea level), average environmental temperature (between 7 and 18 °C), a relative humidity of 80%, and annual rainfall between 800 and 2000 mm. Free grazing was the feeding management system, which was based on mixtures of gramineous and legumes supplemented daily with hay and mineral salts. The cows included were crossbred Holstein and the average farm size was 25.4 ± 4.1 cows, with a milk production between 8 and 12 l/day per cow. In these 15 farms there were 383 cows in production, including 152 multiparous females (3.1 ± 0.17 calvings) with a mean age of 5.9 ± 0.22 years, weight of 553.8 ± 12.82 kg, and body condition of 2.55 ± 0.05 on a scale of 1–5. The animals were an average of 75.3 ± 2.10 d postpartum and in estrus at the time of beginning the study. This requisite was particularly important to enhance the response rate to the synchronization protocol. Of these, finally 140 cows entered the experiment as they responded to the follicular wave restart protocol (described below). The cows were recruited between April 2018 and June 2020.

Animal handling and procedures were performed following the regulations of the Terrestrial Animal Health Code, chapter 7.8 'Use of animals in research and education' of the World Organization for Animal Health (OIE, 2016).

Study design

Cows satisfying the inclusion criteria received the following synchronization protocol to initiate a new follicular wave. On day 1, the cows received an intravaginal progesterone-releasing device (IPD; CIDR® [1.38 g of progesterone], Zoetis, Quito, Ecuador) + 2 mg of intramuscular estradiol benzoate (EB; Gonadiol®, Zoetis). Seven days later, in the morning, an intramuscular dose of 25 mg of prostaglandin (PG; Lutalyse® [dinoprost tromethamine], Zoetis) was applied and the implant was removed. On the following day (day 8), the cows received 1 mg of intramuscular EB (Gonadiol®) and, 54 h after removal of the implant, the location of the preovulatory follicle was explored by transrectal ultrasound (Aloka ProSound 2® Tokyo, Japan) using a 7.5-MHz linear transducer. On day 11, which was considered the day of re-initiation of the follicular wave or 'day 0' of the study, the coccygeal vein was punctured, and blood was sampled into evacuated tubes containing ethylenediaminetetraacetic acid (BD Vacutainer®, Franklin Lakes, NJ, USA). Samples were immediately placed in a cooler with a cooling gel at 5 °C and centrifuged in the laboratory at 1500 g for 20 min. The plasma obtained was stored in aliquots at –20 °C until subsequent analysis of AMH concentrations. On day 11, AFC was also performed (the design is shown in online Supplementary Fig. S1).

AFC

On day 9, the same technician who performed the AFC in all cows also located the preovulatory follicle by transrectal ultrasound. Two days later (day 11), the disappearance of the follicle (ovulation) was verified. In the cows that ovulated, the AFC (<4 mm in diameter) was performed in the two ovaries as described Ayala *et al.* (2019) based on lateral-medial, dorso-ventral, and cranio-caudal scans. The right ovary was scanned first, followed by the left ovary.

Determination of AMH plasma concentration

The Elisa DL-AMH-b kit (DL Develop®, Tech Development, Wuxi, China) was used to measure AMH plasma levels, following the manufacturer's instructions. Optical detection was performed at a wavelength of 450 nm. The manufacturer-reported sensitivity of the assay was <0.117 ng/ml, and the inter- and intra-assay coefficients of variation were <10 and <12%, respectively.

Statistical analysis

Data were stored in Excel and subsequently analyzed using SPSS® 25 (IBM, New York, USA). General descriptive statistics were analyzed. Data were presented in tables as mean \pm standard deviation (SD), while data in the figures were expressed as mean \pm standard error of the mean (SEM). Data that did not follow a normal distribution were \log_{10} -transformed before analysis.

The cows were categorized according to AMH concentration into three groups (low, intermediate, high). Differences in means were assessed for significance using analysis of variance (ANOVA) and compared using the Tukey test with $\alpha = 0.05$. Differences




Fig. 1. Association between plasma levels of anti-Müllerian hormone (AMH) and antral follicle count (AFC) in postpartum cross-bred Holstein cows at the start of a hormonally induced follicular wave.

associated with $P < 0.05$ were considered significant. Pearson's correlation analysis and simple linear regression were applied to explore the potential relationship between AMH levels and AFC.

Results and discussion

AMH plasma levels positively correlated to the total number of antral follicles < 4 mm in diameter present at the restart of the follicular wave ($r = 0.783$; $P < 0.001$). In addition, 61% of the observed variation in average AFC could be explained by the adjusted linear regression model ($r = 0.613$, Fig. 1). Animals presenting plasma AMH concentrations higher than 0.09 ng/ml showed a higher follicular population at the time of restarting the follicular wave than cows with AMH concentrations lower than 0.05 ng/ml. These results confirm the strong relationship between AMH, and the AFC established in the Holstein breed by previous works under normal altitude, observed a correlation with a value of $r = 0.59$ ($P < 0.001$) in cows in stabling (Rico *et al.*, 2009). The correlation was even stronger among bovines raised at sea level ($r = 0.79$, $P < 0.001$; Gobikrushanth *et al.*, 2017).

The linear association observed between plasma AMH and AFC in our work (Fig. 1) is consistent with that described in Holstein cows ($r = 0.61$; $P < 0.001$) and heifers ($r = 0.73$; $P < 0.001$) reared in the low tropics in stabling systems and fed with formulated mixed rations (Batista *et al.*, 2014). Therefore, our data support the belief that AMH plasma concentration can be used as an endocrine indicator to indirectly identify Holstein cows with a higher follicular population at the restart of the new follicular wave under the high-tropic conditions of our study.

Although the association between AMH and AFC in our study is similar to that previously described, the mean value of AMH in our study (0.06 ± 0.03 ng/mL), the minimum value (0.03 ng/ml), and the maximum value (0.16 ng/ml) differed substantially from those previously reported for this breed (online Supplementary Table S1). A study of cows 4–9 years old in the low tropics showed a minimum AMH concentration of 0.025 ng/ml (Rico *et al.*, 2009), similar to the present study. However, the maximum value in that study (0.228 ng/ml) was 1.4 times higher than in our study. The maximum value in our study differed even more from the maximum values in other studies of Holstein cows stabled in the low tropics (0.038–0.774 ng/ml; Gobikrushanth *et al.*, 2017).

These differences in AMH concentrations could reflect the use of different analytical methods. Although all used kits have been validated for use in cattle, they can lead to mean AMH values differing by up to 4.6 times, even if the overall results strongly correlate across the kits ($r = 0.84$, $P < 0.001$) (Fréour *et al.*, 2007). This highlights the relevance of describing the used kits at each study, as well as having previously established cut-off points by each selected diagnostic kit to work with.

Another factor that could affect AMH values is the number of antral follicles < 4 mm present at the time of blood sampling for AMH assessment. In our study, we counted 17.26 ± 0.38 follicles on average (range 7–30), which was lower than the maximum values of 15–55 follicles reported previously (Rico *et al.*, 2009). This lower AFC might explain why our cows showed a lower maximum AMH value. Similarly, other research groups have found lower concentrations of AMH in bovine breeds with a sparse follicular population (Batista *et al.*, 2014). A reduced population of antral follicles has been found in sheep reared at high altitudes, (Parraguez and Gonzalez, 2020). Hypoxia can lead to reduced secretion of follicle-stimulating hormone (FSH) and luteinizing hormone, altering steroidogenesis, and ultimately reducing the number and growth of available follicles. While direct links between hypobaric hypoxia and fertility have not been established in bovines, our work with Creole cattle in the same region as the present study suggest changes in the ovarian physiology, including a lower number of antral follicles (Ayala *et al.*, 2019).

The nutritional conditions typical of the area, characterized by a reduced energy intake, could also influence AFC and AMH levels. Epigenetic studies found that calves born to mothers with nutritional restriction (60% of nutritional requirements during the first 110 d of gestation) presented lower AFC (15.8 ± 1.8 follicles), lower AMH and higher FSH levels than the offspring of mothers that consumed 120% of the nutritional requirements (AFC = 23.6 ± 1.9 ; Mossa *et al.*, 2013). The negative impact of poor maternal nutrition on AFC and AMH levels in the offspring has been confirmed in another study in Holstein cows under intensive production management conditions (Ribeiro *et al.*, 2014). When stratified by AMH levels, 13.6% ($n = 19$) of our cows had high AMH concentrations, 60% ($n = 84$) had intermediate concentrations, and 26.4% ($n = 37$) had low concentrations. The high AMH level (0.14 ± 0.02 ng/ml) was three times the low AMH level (0.04 ± 0.01 ng/ml; $P < 0.05$; Fig. 2a).




Fig. 2. Plasma levels of anti-Müllerian hormone (AMH) (a) and antral follicle count (b) on the day of onset of the follicular wave in cross-bred Holstein cows. Cows were classified into three groups according to the level of AMH, using the cut-off values of 0.09 and 0.05 ng/ml.

Considering that our cattle were exposed to a mixture of different risk factors that have not previously been studied simultaneously, we could not directly infer results on AMH and AFC relationship coming from other studies. The lowest AMH concentration in the present study was similar to that determined previously in Holstein animals with similar parity (Ireland *et al.*, 2011). However, the high concentration group in that study showed 42.8% higher AMH (>0.20 ng/ml) and 58.4% more antral follicles (39.61 ± 2.3 follicles) than our high concentration group. Substantial differences in AFC between animals with 'high' or 'low' AMH were also observed in a study of younger Holsteins (Batista *et al.*, 2014), where the high AMH group had an AMH level of 0.57 ± 0.26 ng/ml and AFC of 34.3 ± 3.12 , while the low AMH group had values of 0.06 ± 0.02 ng/ml and 13.4 ± 1.40 . The lower AFC and AMH in our high AMH animals may reflect the grazing feeding system traditionally provided to herds in the Ecuadorian highlands. The hypobaric hypoxia in the region may then reinforce the effects of this diet on fertility.

Our results suggest that 0.09 ng/ml may be a useful AMH cut-off value for the selection of animals with a sizeable follicular population in the high tropics. With this cut-off, only 13.6% of the cows in our study would be selected, but they would have 1.8 times more antral follicles than the rest. The optimal cut-off point depends on the reproductive technology to be applied. Dairy cows and heifers managed at around sea level, with a follicular population >25 may have the highest number of viable embryos *in vivo*. Animals with a mean AFC of 20.40 ± 1.63 present the best results after ovum pick-up and *in vitro* embryo production (Ghanem *et al.*, 2016). We propose that applying reproductive technologies such as MOET, ovum pick-up and *in vitro* embryo production to crossbred Holstein cows in the high tropics is more likely to be successful if the animals have plasma AMH >0.09 ng/ml and AFC of at least 25.0 ± 2.21 follicles.

In conclusion, crossbred Holstein cows fed on grass living in the Ecuadorian high tropics show a direct relationship between the concentration of AMH and the follicular population. AMH >0.09 ng/ml may be useful for selecting such animals for reproductive biotechnologies. This relationship differed in intensity and shape when compared to those previously observed in other cattle breeds and under different environmental circumstances.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0022029922000140>

Acknowledgements. We thank the farm owners for their support during the experiment. We are also grateful to the veterinarian Ernesto Cobos for his help in sample collection. This work was funded by the Research Directorate of the University of Cuenca (Project No. DIUC_XVII_2018_37_Ayala_Luis).

References

- Alward KJ and Bohlen JF (2020) Overview of anti-Müllerian hormone (AMH) and association with fertility in female cattle. *Reproduction in Domestic Animals* **55**, 3–10.
- Ayala LE, Pesantez JL, Rodas ER, Dután JB, Calle GR, Murillo YA, Vázquez JM, Nieto PE, Ortega VV and Samaniego J (2019) Dinámica folicular de vaquillas Criollas al pastoreo en el altiplano ecuatoriano. *Archivos de Zootecnia* **68**, 186–192.
- Batista EOS, Macedo GG, Sala RV, Ortolan MDDV, Sá Filho MF, Del Valle TA and Baruselli PS (2014) Plasma anti-Müllerian hormone as a predictor of ovarian antral follicular population in *Bos indicus* (Nelore) and *Bos taurus* (Holstein) heifers. *Reproduction in Domestic Animals* **49**, 448–452.
- Fréour T, Miralié S, Bach-Ngohou K, Denis M, Barrière P and Masson D (2007) Measurement of serum anti-Müllerian hormone by Beckman Coulter ELISA and DSL ELISA: comparison and relevance in assisted reproduction technology (ART). *Clinica Chimica Acta* **375**, 162–164.
- Ghanem N, Jin JI, Kim SS, Choi BH, Lee KL, Ha AN and Kong IK (2016) The anti-Müllerian hormone profile is linked with the *in vitro* embryo production capacity and embryo viability after transfer but cannot predict pregnancy outcome. *Reproduction in Domestic Animals* **51**, 301–310.
- Gobikrushanth M, Dutra PA, Bruinjé TC, Colazo MG, Butler ST and Ambrose DJ (2017) Repeatability of antral follicle counts and anti-Müllerian hormone and their associations determined at an unknown stage of follicular growth and an expected day of follicular wave emergence in dairy cows. *Theriogenology* **92**, 90–94.
- Ireland JJA, Smith GWB, Scheetz DA and Folger JKA (2011) Does size matter in females? An overview of the impact of the high variation in the ovarian reserve on ovarian function and fertility, utility of anti-Müllerian hormone as a diagnostic marker for fertility and causes of variation in the ovarian reserve in cattle. *Reproduction, Fertility and Development* **23**, 1–14.
- Jaques JT, Cardoso RC, Chachere J, Sinha S, Seery R, Wilkes AA, Forrest D and Looney CR (2020) Using anti-Müllerian hormone (AMH) as a predictor of ova production for bovine embryo transfer. *Advances in Reproductive Sciences* **8**, 36–47.
- Monniaux D, Rico C, Larroque H, Dalbiès-Tran R, Médigue C, Clément F and Fabre S (2010) L'hormone antimüllérienne, prédicteur endocrinien de la réponse à une stimulation ovarienne chez les bovins. *Gynecologie Obstétrique et Fertilité* **38**, 465–470.
- Mossa F, Carter F, Walsh SW, Kenny DA, Smith GW, Ireland JLH, Hildebrandt TB, Lonergan P, Ireland JJ and Evans ACO (2013) Maternal undernutrition in cows impairs ovarian and cardiovascular systems in their offspring. *Biology of Reproduction* **88**, 1–9.
- Parraguez VH and Gonzalez-Bulnes A (2020) Endocrinology of reproductive function and pregnancy at high altitudes. *Current Opinion in Endocrine and Metabolic Research* **11**, 27–32.
- Perry GA and Cushman RA (2016) Invited review: use of ultrasonography to make reproductive management decisions. *Professional Animal Scientist* **32**, 154–161.

- Ribeiro ES, Bisinotto RS, Lima FS, Greco LF, Morrison A, Kumar A and Santos JEP** (2014) Plasma anti-Müllerian hormone in adult dairy cows and associations with fertility. *Journal of Dairy Science* **97**, 6888–6900.
- Rico C, Fabre S, Medigue C, di Clemente N, Clement F, Bontoux M, Touzé JL, Dupont M, Briant E, Rémy B, Beckers JF and Monniaux D** (2009) Anti-Müllerian hormone is an endocrine marker of ovarian gonadotropin-responsive follicles and can help to predict superovulatory responses in the cow. *Biology of Reproduction* **80**(1), 50–59. <https://doi.org/10.1095/biolreprod.108.072157>.
- Souza AH, Carvalho PD, Rozner AE, Vieira LM, Hackbart KS, Bender RW and Wiltbank MC** (2015) Relationship between circulating anti-Müllerian hormone (AMH) and superovulatory response of high-producing dairy cows. *Journal of Dairy Science* **98**, 169–178.